

A Comparison of different measures for dynamical event-mean transverse momentum fluctuation*

Liu Lianshou¹ and Fu Jinghua^{2,1}

¹*Institute of Particle Physics, Huazhong Normal University, Wuhan 430079 China[†]*

²*Engineering Physics Department, Tsinghua University, Beijing 100084, China*

Various measures for the dynamical event-mean transverse momentum fluctuation are compared with the real dynamical fluctuation using a Monte Carlo model. The variance calculated from the G -moments can reproduce the dynamical variance well, while those obtained through subtraction procedures are approximate measures for not very low multiplicity. The Φ_{p_t} can also serve as an approximate measure after divided by the square root of mean multiplicity.

PACS numbers: 13.85.Hd

Keywords: Multiparticle production; Dynamical fluctuation; Event-by-event analysis

The successful operation of the relativistic heavy ion collider RHIC at BNL has stimulated an intensive study on the theoretically predicted quark gluon plasma (QGP) phase transition. It is expected that the appearance of phase transition will result in an anomalous behavior in the event-by-event fluctuations of transverse momentum [1][2][3].

Since the multiplicity in a single event is finite, there are inevitably statistical fluctuations (SF) in the event-by-event analysis of transverse momentum. Various methods [4][5][6] have been proposed to get rid of the statistical fluctuations and measure the dynamical ones. Most of them are based on a *subtraction* procedure, *i.e.* first to estimate the variance of statistical fluctuations and then subtract it from that of the experimental data. The basic problem in this kind of methods is how to reliably estimate the statistical variance.

Another approach proposed recently [7] is to *eliminate* the SF directly from the experimental data. This method can reproduce the dynamical variance exactly provided the SF are Poissonian, *i.e.* the particles are emitted independently.

The aim of this paper is to compare the different measures proposed in the market and, in particular, to examine the reliability of the estimation of statistical variance. The so called Φ_{p_t} measure [8] will also be discussed and a new measure $\Phi_{p_t}^{(r)}$ will be proposed, which can largely reduce the strong multiplicity dependence of Φ_{p_t} itself.

Let $p(p_t)$ be the dynamical transverse-momentum distribution in single events and

$$p_m = \int_{\Delta} p(p_t) dp_t, \quad (m = 1, 2, \dots, M) \quad (1)$$

be the corresponding “coarse-grained” distribution [9]. The realization of $p(p_t)$ in experiment is the distribution of the total number n of particles in the p_t region Δ , and

the expression

$$q_m = n_m/n \quad (m = 1, 2, \dots, M) \quad (2)$$

is an evaluation of p_m , where n_m is the number of particles falling into the m th bin. Thus the dynamical and experimentally-measured event-mean transverse momentum are, respectively,

$$\bar{p}_{t\text{dyn}} = \int_{\Delta} p_t p(p_t) dp_t = \sum_{m=1}^M (p_t)_m p_m, \quad (3)$$

$$\bar{p}_{t\text{exp}} = \sum_{m=1}^M (p_t)_m q_m = \sum_{m=1}^M (p_t)_m \frac{n_m}{n}, \quad (4)$$

where $(p_t)_m$ is the p_t value in the m th bin. The event-space moments of $\bar{p}_{t\text{dyn}}$ and $\bar{p}_{t\text{exp}}$ are

$$C_p^{\text{dyn}}(\bar{p}_t) = \langle \bar{p}_{t\text{dyn}}^p \rangle = \left\langle \left(\sum_{m=1}^M (p_t)_m p_m \right)^p \right\rangle, \quad (5)$$

$$C_p^{\text{exp}}(\bar{p}_t) = \langle (\bar{p}_{t\text{exp}})^p \rangle = \left\langle \left(\sum_{m=1}^M (p_t)_m \frac{n_m}{n} \right)^p \right\rangle, \quad (6)$$

respectively. These two are evidently unequal due to the existence of SF. In particular, the experimental moments $C_p^{\text{exp}}(\bar{p}_t)$ depend crucially on the particle multiplicity [11], which is mainly due to SF. Therefore, subtraction procedures are commonly used to extract the dynamical fluctuation of $\bar{p}_t$.

The basic idea of the subtraction procedure is that the variances $\sigma^2(\bar{p}_t) = C_2(\bar{p}_t) - C_1(\bar{p}_t)^2$ of dynamical and statistical fluctuations are additive

$$\sigma_{\bar{p}_t\text{data}}^2 = \sigma_{\bar{p}_t\text{dyn}}^2 + \sigma_{\bar{p}_t\text{stat}}^2, \quad (7)$$

where $\sigma_{\bar{p}_t\text{data}}^2$, $\sigma_{\bar{p}_t\text{dyn}}^2$ and $\sigma_{\bar{p}_t\text{stat}}^2$ are the experimental, dynamical and statistical variances, respectively. This additivity is true if the DF and SF are independent, which

*This work is supported in part by the NSFC under project 90103019.

[†]Electronic address: liuls@iopp.ccnu.edu.cn

is a reasonable assumption. Therefore, the main problem in the subtraction procedure for extracting the dynamical variance is to estimate the statistical ones. Various methods have been proposed for this purpose.

For example, in Ref. [4] the results from mixed events are considered as the baseline for the random distribution and the difference in the fluctuation from a random distribution defined as

$$d = \omega_{\text{data}} - \omega_{\text{baseline}} \quad (8)$$

is taken as a measure of the dynamical fluctuation. In Eq.(8)

$$\omega = \frac{\sqrt{\langle \bar{p}_t^2 \rangle - \langle \bar{p}_t \rangle^2}}{\langle \bar{p}_t \rangle} = \frac{\sqrt{\sigma_{\bar{p}_t}^2}}{\langle \bar{p}_t \rangle}, \quad (9)$$

$\bar{p}_t$ is the mean transverse momentum in a single event and $\langle \dots \rangle$ denotes the average over event sample.

Alternatively, in Ref. [5] the statistical variance of event mean p_t , under the assumption of independent particle production, is estimated as

$$\sigma_{\bar{p}_t \text{stat}}^2 = \frac{\sigma_{p_t \text{incl}}^2}{\langle n \rangle}, \quad (10)$$

and, therefore, the dynamical variance is equal to

$$\sigma_{\bar{p}_t \text{VKR}}^2 = \sigma_{\bar{p}_t \text{data}}^2 - \sigma_{\bar{p}_t \text{stat}}^2 = \sigma_{\bar{p}_t \text{data}}^2 - \frac{\sigma_{p_t \text{incl}}^2}{\langle n \rangle}. \quad (11)$$

A somewhat different expression is used in Ref. [6], where the experimental variance is weighted by event multiplicity N_j :

$$\sigma_{\bar{p}_t \text{CERES}}^2 = \frac{\sum_{j=1}^{\mathcal{N}} N_j (\bar{p}_t^j - \langle \bar{p}_t \rangle)^2}{\sum_{j=1}^{\mathcal{N}} N_j} - \frac{\sigma_{p_t \text{incl}}^2}{\langle n \rangle}, \quad (12)$$

where $\mathcal{N}$ is the total number of events.

A widely used measure for the non-statistical mean p_t fluctuation is the Φ_{p_t} proposed in Ref. [8]

$$\Phi_{p_t} \equiv \sqrt{\langle Z^2 \rangle / \langle n \rangle} - \sqrt{\langle z^2 \rangle}, \quad (13)$$

where z and Z are defined as $z \equiv p_t - \langle p_t \rangle$ for each particle and $Z \equiv \sum_{i=1}^n z_i = n(\bar{p}_t - \langle p_t \rangle)$ for each event, respectively [10]. The second term in the r.h.s. of Eq. (13) is the square root of the inclusive variance $\sigma_{p_t \text{incl}}^2 = \langle (p_t - \langle p_t \rangle)^2 \rangle$. Assuming that the multiplicity fluctuation is uncorrelated with the p_t fluctuation, we get from Eq.(13)

$$\Phi_{p_t} = \sqrt{\langle n^2 \rangle / \langle n \rangle} \sigma_{\bar{p}_t \text{data}} - \sigma_{p_t \text{incl}}. \quad (14)$$

This equation is evidently similar to Eq.(11).

All of the above measures have the same structure, being based on a subtraction procedure, *i.e.* to subtract the variance of $\bar{p}_t$ or a quantity related to it, that will be expected from a pure statistical system, from the same




FIG. 1: The experimental (open triangles) and dynamical (dashed line) variances of $\bar{p}_t$, together with the variances calculated from G -moments (open circles) and from Eqs. (11) (upward solid triangles) and (12) (downward solid triangles).

quantity obtained in experiment. These measures will, of course, vanish for a pure statistical system, and a non-vanishing value of them will indicate the existence of dynamical effect. Therefore, the measures based on the subtraction procedure, as those listed above, will at least qualitatively measure the effect of DF.

In order to have an idea on how the various subtraction methods work quantitatively, let us carry on a Monte Carlo simulation using a toy model. In this model the distribution of p_t is taken as

$$p(p_t) = \frac{4}{a^2} p_t e^{-2p_t/a}, \quad (15)$$

with a Gaussian distributed parameter a ($\sigma^2(a) = 0.24$). In total $20 \times 500,000$ events are generated for 20 average multiplicities. The resulting dynamical and experimental variances $\sigma_{\bar{p}_t \text{dyn}}^2$ and $\sigma_{\bar{p}_t \text{exp}}^2$ are plotted in Fig.1 as dashed line and open triangles, respectively. For comparison, the $\sigma_{\bar{p}_t \text{VKR}}^2$ and $\sigma_{\bar{p}_t \text{CERES}}^2$ are also plotted in the same figure as upward and downward solid triangles, respectively.

The differences between the experimental and dynamical variances $\sigma_{\bar{p}_t \text{exp}}^2 - \sigma_{\bar{p}_t \text{dyn}}^2$ are due to SF. It can be seen from the figure that the $\sigma_{\bar{p}_t \text{VKR}}^2$ indeed reduces the SF, especially when the average multiplicity is not very low; while the $\sigma_{\bar{p}_t \text{CERES}}^2$ overshoots the influence of SF.

On the other hand, the open circles shown in the figure reproduce the dynamical variances very well. They are obtained through a totally different approach [7], which is based on the elimination of the SF directly from the experimental data.

For the first order moment the elimination of SF is straightforward,

$$C_1^{(\text{dyn})}(\bar{p}_t) = G_1(\bar{p}_t) = \left\langle \sum_{m=1}^M (p_t)_m \frac{n_m}{\langle n \rangle} \right\rangle, \quad (16)$$

provided the SF is Poissonian. The key point in eliminating SF in the second order moment is to expand the 2nd power in the definition Eq.(5) of $C_2^{(\text{dyn})}(\bar{p}_t)$,

$$C_2^{(\text{dyn})}(\bar{p}_t) = \left\langle \left(\sum_{m=1}^M (p_t)_m p_m \right)^2 \right\rangle = \left\langle \sum_{m=1}^M (p_t)_m^2 p_m^2 \right\rangle + \left\langle \sum_{m \neq m'}^M (p_t)_m (p_t)_{m'} p_m p_{m'} \right\rangle. \quad (17)$$

Using the formulae

$$\left\langle \sum_{m=1}^M f_m p_m^p \right\rangle = \left\langle \sum_{m=1}^M f_m \frac{n_m(n_m-1) \cdots (n_m-p+1)}{\langle n \rangle^p} \right\rangle, \quad (18)$$

which holds for Poissonian SF, we then get $C_2^{(\text{dyn})}(\bar{p}_t) = G_2(\bar{p}_t)$, where

$$G_2(\bar{p}_t) = \left\langle \sum_{m=1}^M (p_t)_m^2 \frac{n_m(n_m-1)}{\langle n \rangle^2} \right\rangle + \left\langle \sum_{m \neq m'}^M (p_t)_m (p_t)_{m'} \frac{n_m n_{m'}}{\langle n \rangle^2} \right\rangle. \quad (19)$$




FIG. 2: Comparison of $\sigma_{\bar{p}_t \text{stat}}^2$ and $\sigma_{\bar{p}_t \text{incl}}^2 / \langle n \rangle$ for systems with and without DF

This guarantees the equality of G -variance $\sigma_{\bar{p}_t G}^2 = G_2(\bar{p}_t) - G_1(\bar{p}_t)^2$ to dynamical ones $\sigma_{\bar{p}_t \text{dyn}}^2 = C_2^{(\text{dyn})}(\bar{p}_t) - C_1^{(\text{dyn})}(\bar{p}_t)^2$.

On the contrary, the SF cannot be exactly gotten rid of using the subtraction procedure. The main problem lies in the estimation of statistical variance by Eq.(10). This

equality holds only for a pure statistical system without any dynamical fluctuation. When dynamical fluctuations are present the $\sigma_{\bar{p}_t \text{stat}}^2$ and $\sigma_{\bar{p}_t \text{incl}}^2 / \langle n \rangle$ are, in general, unequal. For an exhibition, the $\sigma_{\bar{p}_t \text{stat}}^2 = \sigma_{\bar{p}_t \text{exp}}^2 - \sigma_{\bar{p}_t \text{dyn}}^2$, cf. Fig.1, and $\sigma_{\bar{p}_t \text{incl}}^2 / \langle n \rangle$ are plotted in Fig.2 as solid circles and solid triangles, respectively. They are approximately equal to each other only when the average multiplicity is not very low, *e.g.* when $\langle n \rangle > 12$.

In order to further clarify this point, we have made a pure statistical model with $p_m = 1/M$, cf. Eq.1. In this case $\sigma_{\bar{p}_t \text{dyn}}^2 \equiv 0$. The corresponding $\sigma_{\bar{p}_t \text{stat}}^2 = \sigma_{\bar{p}_t \text{exp}}^2$ and $\sigma_{\bar{p}_t \text{incl}}^2 / \langle n \rangle$ are plotted in Fig.2 as open circles and open triangles, respectively. The differences between them are small up to $\langle n \rangle = 5$. This shows that the discrepancy between $\sigma_{\bar{p}_t \text{VKR}}^2$ ($\sigma_{\bar{p}_t \text{CERES}}^2$) and $\sigma_{\bar{p}_t \text{dyn}}^2$ is mainly due to the inappropriate estimation at low multiplicity of $\sigma_{\bar{p}_t \text{stat}}^2$ by $\sigma_{\bar{p}_t \text{incl}}^2 / \langle n \rangle$ for a system containing DF.

Let us now turn to the discussion on the Φ_{p_t} measure [8][11]. The Monte Carlo results of Φ_{p_t} are plotted in Fig.3 as solid circles. It is strongly multiplicity dependent, and is, therefore, not a good measure for DF. On accounting of Eq.(14) we divide Φ_{p_t} by the square root of average multiplicity and define

$$\Phi_{p_t}^{(r)} \equiv \frac{\Phi_{p_t}}{\sqrt{\langle n \rangle}} \approx \sigma_{\bar{p}_t \text{data}} - \frac{\sigma_{\bar{p}_t \text{incl}}}{\sqrt{\langle n \rangle}}, \quad (20)$$

cf. Eq.(11).

The Monte Carlo results of $\Phi_{p_t}^{(r)}$ are shown in Fig.3 as upward solid triangles together with $\sigma_{\bar{p}_t \text{CERES}} = \sqrt{\sigma_{\bar{p}_t \text{CERES}}^2}$ (downward triangles) and $\sigma_{\bar{p}_t \text{dyn}}$ (dashed




FIG. 3: Variation of Φ_{pt} and Φ_{pt}^r with $\langle n \rangle$, compared with $\sigma_{\bar{p}_t}^{CERES}$ and $\sigma_{\bar{p}_t}^{dyn}$

line). It can be seen from the figure that $\Phi_{pt}^{(r)}$ has a much

weaker multiplicity dependence and is closer to $\sigma_{\bar{p}_t}^{dyn}$ than Φ_{pt} . Therefore, $\Phi_{pt}^{(r)}$ is a better measure of DF than Φ_{pt} itself.

In this paper, various measures for the dynamical transverse momentum fluctuation in event-by-event analysis are compared with the dynamical fluctuation using a Monte Carlo model.

It turns out that the $\sigma_{\bar{p}_t}^2$ calculated from the G-moments coincides with the dynamical variance $\sigma_{\bar{p}_t}^{2,dyn}$, showing that the G-moment method is effective in eliminating the statistical fluctuations coming from the finite number of particle in a single event and thus provides a good measure for the dynamical fluctuations.

The $\sigma_{\bar{p}_t}^2$ proposed by S. A. Voloshin, V. Koch and H. G. Ritter [5] and its revised version $\sigma_{\bar{p}_t}^2$ [6] are found to be good approximate measures for the dynamical fluctuation when the multiplicity is not very low. Therefore, their application to relativistic heavy ion experiments is justified.

The Φ_{pt} divided by the square root of average multiplicity ($\Phi_{pt}^r = \Phi_{pt}/\sqrt{\langle n \rangle}$) is another approximate measure for the dynamical fluctuation, which depends on multiplicity weaker than Φ_{pt} and is closer to the square root of dynamical variance $\sigma_{\bar{p}_t}^{dyn}$.

Acknowledgement The authors thank Wu Yuanfang and Liu Feng for helpful discussions and comments.

-
- [1] S. Mrówczyński, Phys. Lett. **B314** (1993) 118.
 - [2] M. A. Stephanov, K. Rajagopal and E. V. Shuryak, Phys. Rev. Lett. **81** (1998) 4816; Phys. Rev. **D60** (1999) 114028.
 - [3] A. Dumitru and R. Pisarski, Phys. Lett. **B504** (2001) 282.
 - [4] K. Adcox *et al.* (PHENIX Collab.), nucl-ex/0203015 (2002).
 - [5] S. A. Voloshin, V. Koch and H. G. Ritter, Phys. Rev. **C60**, 024901 (1999).
 - [6] D. Adamová *et al.* (CERES Collab.), nucl-ex/0305002 (2003).
 - [7] Fu Jinghua and Liu Lianshou, *Extracting Event Dynamics from Event-by-Event Analysis* Phys. Rev. **C**, to be published.
 - [8] M. Gaździcki and S. Mrówczyński, Z. Phys. **C54**, 127 (1992); F. Liu, A. Tai M. Gaździcki and R. Stock, Eur. Phys. J. **C8**, 649 (1999).
 - [9] A. Białas and R. Peschanski, Nucl. Phys. **B273**, 703 (1986); *ibid* **308**, 857 (1988).
 - [10] Through out the present paper the symbol $\langle x \rangle$ is used to denote the sample average of x , no matter x is an event variable or an inclusive variable, and the symbol $\bar{x}$ is reserved for the event average of x .
 - [11] T. Anticic *et al.* (NA49 Collaboration), hep-ex/0311990 (2003).